

Generalized Enrichment of Categories

Tom Leinster*

Department of Pure Mathematics and Mathematical Statistics,
Centre for Mathematical Sciences, Wilberforce Road,
Cambridge CB3 0WB, UK
`leinster@dpms.cam.ac.uk`

Abstract

We define the phrase ‘category enriched in an **fc**-multicategory’ and explore some examples. An **fc**-multicategory is a very general kind of 2-dimensional structure, special cases of which are double categories, bicategories, monoidal categories and ordinary multicategories. Enrichment in an **fc**-multicategory extends the (more or less well-known) theories of enrichment in a monoidal category, in a bicategory, and in a multicategory. Moreover, **fc**-multicategories provide a natural setting for the bimodules construction, traditionally performed on suitably cocomplete bicategories. Although this paper is elementary and self-contained, we also explain why, from one point of view, **fc**-multicategories are the *natural* structures in which to enrich categories.

Published as *Journal of Pure and Applied Algebra* **168** (2002), 391–406.

2000 Mathematics Subject Classification: 18D20, 18D05, 18D50, 18D10.

A general question in category theory is: given some kind of categorical structure, what might it be enriched in? For instance, suppose we take braided monoidal categories. Then the question asks: what kind of thing must $\mathcal{V}$ be if we are to speak sensibly of $\mathcal{V}$ -enriched braided monoidal categories? (The usual answer is that $\mathcal{V}$ must be a symmetric monoidal category.)

In another paper, [7], I have given an answer to the general question for a certain family of categorical structures (generalized multicategories). In particular, this theory gives an answer to the question ‘what kind of structure $\mathcal{V}$ can a category be enriched in’? The answer is: an ‘**fc**-multicategory’.

Of course, the traditional answer to this question is that $\mathcal{V}$ is a monoidal category. But there is also a notion of a category enriched in a bicategory (see Walters [15]). And generalizing in a different direction, it is easy to see how

*Supported by the EPSRC and St John’s College, Cambridge

one might speak of a category enriched in an ordinary multicategory (‘change tensors to commas’). An **fc**-multicategory is, in fact, a very general kind of 2-dimensional categorical structure, encompassing monoidal categories, bicategories, multicategories and double categories. The theory of categories enriched in an **fc**-multicategory extends all of the aforementioned theories of enrichment.

So from the point of view of [7], **fc**-multicategories are the natural structures in which to enrich a category. In this work, however, we do not assume any knowledge of [7] or of generalized multicategories. Instead, we define **fc**-multicategory in an elementary fashion (Section 1) and then define what a category enriched in an **fc**-multicategory is (Section 2). Along the way we see how enrichment in an **fc**-multicategory extends the previously-mentioned theories of enrichment, and look at various examples.

fc-multicategories also provide a natural setting for the bimodules construction (Section 3), traditionally carried out on bicategories satisfying certain co-completeness conditions. At the level of **fc**-multicategories, the construction is both more general and free of technical restrictions. We show, in particular, that a category enriched in an **fc**-multicategory $\mathcal{V}$ naturally gives rise to a category enriched in the **fc**-multicategory $\mathbf{Bim}(\mathcal{V})$ of bimodules in $\mathcal{V}$. This result is functorial (that is, a $\mathcal{V}$ -enriched functor gives rise to a $\mathbf{Bim}(\mathcal{V})$ -enriched functor), a statement which only holds if we work with **fc**-multicategories rather than bicategories.

1 **fc**-multicategories

In a moment, an explicit and elementary definition of **fc**-multicategory will be given. But first it might be helpful to look briefly at the wider context in which this definition sits: the theory of ‘generalized multicategories’. The reader is reassured that no knowledge of this wider context is required in order to understand the rest of the paper.

Given a monad T on a category $\mathcal{E}$, both having certain convenient properties, there is a category of T -multicategories. A T -multicategory C consists of a diagram

$$\begin{array}{ccc} & C_1 & \\ \text{dom} \swarrow & & \searrow \text{cod} \\ T(C_0) & & C_0 \end{array}$$

in $\mathcal{E}$ (a T -graph) together with functions defining ‘composition’ and ‘identity’; the full details can be found in Burroni [3] or Leinster ([6] or [8]). Thus when T is the identity monad on $\mathcal{E} = \mathbf{Set}$, a T -multicategory is simply a category. When T is the free-monoid monad on $\mathcal{E} = \mathbf{Set}$, a T -multicategory is a multicategory in the original sense of Lambek [5]. When T is the free (strict) ∞ -category monad on the category $\mathcal{E}$ of globular sets (‘ ∞ -graphs’), a T -multicategory C with $C_0 = 1$ is a higher operad in the sense of Batanin [1]. The example which concerns us here is when T is the free category monad **fc** on the category $\mathcal{E}$ of

directed graphs. A T -multicategory is then an **fc**-multicategory in the sense of the following explicit definition.

Definition 1 *An **fc**-multicategory consists of*

- A class of objects $x, x', \dots$
- For each pair (x, x') of objects, a class of vertical 1-cells $\begin{array}{c} x \\ \downarrow \\ x' \end{array}$, denoted $f, f', \dots$
- For each pair (x, x') of objects, a class of horizontal 1-cells $x \longrightarrow x'$, denoted $m, m', \dots$
- For each $n \geq 0$, objects $x_0, \dots, x_n, x, x'$, vertical 1-cells f, f' , and horizontal 1-cells $m_1, \dots, m_n, m$, a class of 2-cells

$$\begin{array}{ccccccc} x_0 & \xrightarrow{m_1} & x_1 & \xrightarrow{m_2} & \dots & \xrightarrow{m_n} & x_n \\ f \downarrow & & & & \Downarrow & & \downarrow f' \\ x & \xrightarrow{\quad m \quad} & & & & & x' \end{array} \quad (1)$$

denoted $\theta, \theta', \dots$

- Composition and identity functions making the objects and vertical 1-cells into a category
- A composition function for 2-cells, as in the picture

$$\begin{array}{ccccccc} \bullet & \xrightarrow{m_1^1} & \dots & \xrightarrow{m_1^{r_1}} & \bullet & \xrightarrow{m_2^1} & \dots & \xrightarrow{m_2^{r_2}} & \bullet & \dots & \bullet & \xrightarrow{m_n^1} & \dots & \xrightarrow{m_n^{r_n}} & \bullet \\ f_0 \downarrow & & \Downarrow \theta_1 & & \downarrow & & \Downarrow \theta_2 & & \downarrow & \dots & \downarrow & & \Downarrow \theta_n & & \downarrow f_n \\ \bullet & \xrightarrow{\quad m_1 \quad} & \bullet & \xrightarrow{\quad m_2 \quad} & \bullet & \dots & \bullet & \xrightarrow{\quad m_n \quad} & \bullet \\ f \downarrow & & & & \Downarrow \theta & & & & \downarrow f' \\ \bullet & \xrightarrow{\quad m \quad} & \bullet \end{array}$$

$\longmapsto$

$$\begin{array}{ccc} \bullet & \xrightarrow{m_1^1} & \dots & \xrightarrow{m_n^{r_n}} & \bullet \\ f \circ f_0 \downarrow & & \Downarrow \theta \circ (\theta_1, \theta_2, \dots, \theta_n) & & \downarrow f' \circ f_n \\ \bullet & \xrightarrow{\quad m \quad} & \bullet \end{array}$$

($n \geq 0, r_i \geq 0$), where the $\bullet$'s represent objects

- An identity function

$$x \xrightarrow{m} x' \quad \longmapsto \quad \begin{array}{ccc} x & \xrightarrow{m} & x' \\ 1_x \downarrow & \Downarrow 1_m & \downarrow 1_{x'} \\ x & \xrightarrow{m} & x' \end{array}$$

The 2-cell composition and identities are required to obey associativity and identity laws.

The associativity and identity laws ensure that any diagram of pasted-together 2-cells with a rectangular boundary has a well-defined composite.

Examples

1. Any double category gives an **fc**-multicategory, in which a 2-cell as in diagram (1) is a 2-cell

$$\begin{array}{ccc} x_0 & \xrightarrow{m_n \circ \dots \circ m_1} & x_n \\ f \downarrow & \Downarrow & \downarrow f' \\ x & \xrightarrow{m} & x' \end{array}$$

in the double category. If the double category is called $\mathcal{D}$ then we also call the resulting **fc**-multicategory $\mathcal{D}$, and we use the same convention for bicategories (next example).

2. Any bicategory gives an **fc**-multicategory in which the only vertical 1-cells are identity maps, and a 2-cell as in diagram (1) is a 2-cell

$$\begin{array}{ccc} & m_n \circ \dots \circ m_1 & \\ x_0 & \Downarrow & x_n \\ & m & \end{array}$$

in the bicategory (with $x = x_0$ and $x' = x_n$).


Here $m_n \circ \dots \circ m_1$ denotes some n -fold composite of the 1-cells $m_n, \dots, m_1$ in the bicategory. For the sake of argument let us decide to associate to the left, so that $m_4 \circ m_3 \circ m_2 \circ m_1$ means $((m_4 \circ m_3) \circ m_2) \circ m_1$. A different choice of bracketing would only affect the resulting **fc**-multicategory up to isomorphism (in the obvious sense).

3. Any monoidal category $\mathcal{M}$ gives rise to an **fc**-multicategory $\Sigma\mathcal{M}$ (the *suspension* of $\mathcal{M}$) in which there is one object and one vertical 1-cell, and a 2-cell


$$\begin{array}{ccc} \bullet & \xrightarrow{M_1} \bullet & \xrightarrow{M_2} \bullet & \dots & \bullet & \xrightarrow{M_n} \bullet \\ 1 \downarrow & & & & & \downarrow 1 \\ \bullet & \xrightarrow{M} & \bullet & & & \bullet \end{array} \quad (2)$$

is a morphism $M_n \otimes \dots \otimes M_1 \longrightarrow M$ in $\mathcal{M}$. This is a special case of Example (2).


4. Similarly, any ordinary multicategory $\mathcal{M}$ gives an **fc**-multicategory $\Sigma\mathcal{M}$: there is one object, one vertical 1-cell, and a 2-cell as in diagram (2) is a map $M_1, \dots, M_n \longrightarrow M$ in $\mathcal{M}$.
5. We define an **fc**-multicategory **Span**. Objects are sets, vertical 1-cells are functions, a horizontal 1-cell $X \longrightarrow Y$ is a diagram



of sets and functions, and a 2-cell inside



is a function θ making



commute. Here $M_n \circ \dots \circ M_1$ is the limit of the top row of diagram (3), an iterated pullback. Composition is defined in the obvious way.

Span is an example of a ‘weak double category’, which is just like a double category except that horizontal 1-cell composition only obeys associativity and identity axioms up to coherent isomorphism.

It is rather idiosyncratic to name this **fc**-multicategory after its horizontal 1-cells: usually one names a categorical structure after its objects (e.g. **Group**, **Set**). However, we do not want to confuse the **fc**-multicategory **Span** of sets with the mere category **Set** of sets, so we will stick to this convention.

Notice, incidentally, that **Set** is the category formed by the objects and vertical 1-cells of **Span**, and that the **fc**-multicategory $\Sigma\mathbf{Set}$ arising from the monoidal category $(\mathbf{Set}, \times, 1)$ is the ‘full’ sub-**fc**-multicategory of **Span** whose only object is 1.

6. There is an **fc**-multicategory **Prof**, in which the category formed by the objects and vertical 1-cells is the usual category of (small) categories and

functors. Horizontal 1-cells are profunctors (bimodules): that is, a horizontal 1-cell $\mathbb{X} \longrightarrow \mathbb{X}'$ is a functor $\mathbb{X}^{\text{op}} \times \mathbb{X}' \longrightarrow \mathbf{Set}$. A 2-cell

$$\begin{array}{ccccccc} \mathbb{X}_0 & \xrightarrow{M_1} & \mathbb{X}_1 & \xrightarrow{M_2} & \cdots & \xrightarrow{M_n} & \mathbb{X}_n \\ F \downarrow & & & & \Downarrow & & \downarrow F' \\ \mathbb{X} & \xrightarrow{\quad M \quad} & & & & & \mathbb{X}' \end{array}$$

consists of a function

$$M_n(x_{n-1}, x_n) \times \cdots \times M_1(x_0, x_1) \longrightarrow M(F(x_0), F'(x_n))$$

for each $x_0 \in \mathbb{X}_0, \dots, x_n \in \mathbb{X}_n$, such that this family of functions is natural in the x_i 's. So if the functors F and F' are identities then this is a morphism of profunctors $M_n \otimes \cdots \otimes M_1 \longrightarrow M$.

7. In a similar spirit, **Bimod** is the following **fc**-multicategory:

- objects are rings (with identity, not necessarily commutative)
- vertical 1-cells are ring homomorphisms
- a horizontal 1-cell $R \longrightarrow S$ is an (S, R) -bimodule
- a 2-cell

$$\begin{array}{ccccccc} R_0 & \xrightarrow{M_1} & R_1 & \xrightarrow{M_2} & \cdots & \xrightarrow{M_n} & R_n \\ f \downarrow & & & & \Downarrow \theta & & \downarrow f' \\ R & \xrightarrow{\quad M \quad} & & & & & R' \end{array}$$

is a function $\theta : M_n \times \cdots \times M_1 \longrightarrow M$ which preserves addition in each component separately (is ‘multi-additive’) and satisfies the equations

$$\begin{aligned} \theta(r_n \cdot m_n, m_{n-1}, \dots) &= f'(r_n) \cdot \theta(m_n, m_{n-1}, \dots) \\ \theta(m_n \cdot r_{n-1}, m_{n-1}, \dots) &= \theta(m_n, r_{n-1} \cdot m_{n-1}, \dots) \\ &\vdots \\ \theta(\dots, m_2 \cdot r_1, m_1) &= \theta(\dots, m_2, r_1 \cdot m_1) \\ \theta(\dots, m_2, m_1 \cdot r_0) &= \theta(\dots, m_2, m_1) \cdot f(r_0) \end{aligned}$$

- composition and identities are defined in the evident way.

8. If we remove all the additive structure involved in **Bimod** then we obtain an **fc**-multicategory **Action**; alternatively, **Action** is the ‘full’ sub-**fc**-multicategory of **Prof** in which the only objects allowed are 1-object categories. Thus the objects of **Action** are monoids, the vertical 1-cells are monoid homomorphisms, a horizontal 1-cell $R \longrightarrow S$ is a set with commuting left S -action and right R -action, and 2-cells are defined as in Example (7).

2 Enrichment

The purpose of this paper is to explore in an elementary way the concept of a category enriched in an **fc**-multicategory. But just as the elementary definition of **fc**-multicategory (Definition 1) is plucked out of a much larger theory (as explained in the introduction to Section 1), so too is the definition of category enriched in an **fc**-multicategory. There is a whole theory [7] of enrichment for generalized multicategories, of which the present work is just the most simple case. This wider theory runs as follows.

Any T -multicategory has an underlying T -graph, as explained above, and so there is a forgetful functor

$$T\text{-Multicat} \longrightarrow T\text{-Graph}.$$

Under mild conditions on $\mathcal{E}$ and T , this functor has a left adjoint. We thus obtain a monad T' on the category $\mathcal{E}' = T\text{-Graph}$. We can then speak of T' -multicategories, and if $\mathcal{V}$ is a T' -multicategory one can make a definition of $\mathcal{V}$ -enriched T -multicategory. So: we can speak of a T -multicategory enriched in a T' -multicategory.

The most simple case is the identity monad T on $\mathcal{E} = \mathbf{Set}$. Then T -multicategories are categories, T' is the free category monad **fc** on $\mathcal{E}' = \mathbf{Graph}$, and T' -multicategories are **fc**-multicategories. So the general theory gives a concept of category enriched in an **fc**-multicategory. The main part of this section is a direct description of this concept.

The next most simple case is the free monoid monad T on $\mathcal{E} = \mathbf{Set}$, and here there are two especially interesting examples of enriched T -multicategories. Firstly, it turns out that any symmetric monoidal category $\mathcal{S}$ gives rise to a T' -multicategory $\mathcal{V}$, and a one-object $\mathcal{V}$ -enriched T -multicategory is then exactly what topologists call a (non-symmetric) operad in $\mathcal{S}$ (see e.g. [9]). Secondly, there is a certain naturally-arising T' -multicategory $\mathcal{V}$ such that $\mathcal{V}$ -enriched T -multicategories are the structures called ‘relaxed multicategories’ by Borchers in his definition of vertex algebras over a vertex group ([2], [10], [11]), and called ‘pseudo-monoidal categories’ by Soibelman in his work on quantum affine algebras ([12], [13]).

The general definition of enriched T -multicategory is very simple. Take a monad T on a category $\mathcal{E}$, and let T' be the free T -multicategory monad, as above. Given an object A of $\mathcal{E}$, we can form $I(A)$ (with I for ‘indiscrete’), the unique T -multicategory with graph

$$T(A) \xleftarrow{\text{pr}_1} T(A) \times A \xrightarrow{\text{pr}_2} A.$$

Then $I(A)$ is a T' -algebra, say $h : T'(I(A)) \longrightarrow I(A)$. Arising from this is a T' -multicategory $M(I(A))$, the unique such with graph

$$T'(I(A)) \xleftarrow{1} T'(I(A)) \xrightarrow{h} I(A).$$

For a fixed T' -multicategory $\mathcal{V}$, a $\mathcal{V}$ -enriched T -multicategory is defined as an object C_0 of $\mathcal{E}$ together with a map $T'(I(C_0)) \longrightarrow \mathcal{V}$ of T' -multicategories.

Maps between $\mathcal{V}$ -enriched T -multicategories are also defined in a simple way, thus giving a category.

In the case concerning us, $\mathcal{E} = \mathbf{Set}$ and $T = id$, the definition of enriched (T -multi)category is therefore as follows. Given a set A , we obtain the indiscrete category $I(A)$ on A . In the $\mathbf{fc}$ -multicategory $M(I(A))$, an object is an element of A , the only vertical 1-cells are identities, there is one horizontal 1-cell $a \longrightarrow b$ for each $a, b \in A$, and for each $a_0, \dots, a_n \in A$ there is precisely one 2-cell of the form

$$\begin{array}{ccccccc} a_0 & \longrightarrow & a_1 & \longrightarrow & \cdots & \longrightarrow & a_n \\ \downarrow 1 & & & & \Downarrow & & \downarrow 1 \\ a_0 & \longrightarrow & & & & & a_n \end{array}$$

Composition and identities are uniquely determined. A category enriched in an $\mathbf{fc}$ -multicategory $\mathcal{V}$ consists of a set C_0 together with a map from the $\mathbf{fc}$ -multicategory $M(I(C_0))$ to $\mathcal{V}$. This definition is plainly equivalent to Definition 2 below.

That concludes the sketch of the theory of enriched generalized multicategories, and we now return to the elementary account.

Fix an $\mathbf{fc}$ -multicategory $\mathcal{V}$.

Definition 2 A category enriched in $\mathcal{V}$, or $\mathcal{V}$ -enriched category, C , consists of

- a class C_0 (of ‘objects’)
- for each $a \in C_0$, an object $C[a]$ of $\mathcal{V}$
- for each $a, b \in C_0$, a horizontal 1-cell $C[a] \xrightarrow{C[a,b]} C[b]$ in $\mathcal{V}$
- for each $a, b, c \in C_0$, a ‘composition’ 2-cell

$$\begin{array}{ccccc} C[a] & \xrightarrow{C[a,b]} & C[b] & \xrightarrow{C[b,c]} & C[c] \\ \downarrow 1 & & \Downarrow \text{comp}_{a,b,c} & & \downarrow 1 \\ C[a] & \xrightarrow{C[a,c]} & & & C[c] \end{array}$$

- for each $a \in C_0$, an ‘identity’ 2-cell

$$\begin{array}{ccc} C[a] & \xlongequal{\quad} & C[a] \\ \downarrow 1 & \Downarrow \text{ids}_a & \downarrow 1 \\ C[a] & \xrightarrow{C[a,a]} & C[a] \end{array}$$

(where the equality sign along the top denotes a string of 0 horizontal 1-cells)

such that *comp* and *ids* satisfy associativity and identity axioms.

To the reader used to enrichment in a monoidal category, the only unfamiliar piece of data in this definition is the family of objects $C[a]$. To the reader used to enrichment in bicategories even this will be familiar; indeed, since the vertical 1-cells are not used in any significant way, our definition looks very much like the definition of category enriched in a bicategory (see [15]). This lack of use of the vertical 1-cells might seem to weigh against the claim that **fc**-multicategories are, in some sense, the natural structures in which to enrich categories. However, the vertical 1-cells *are* used in the definition of $\mathcal{V}$ -enriched functor, which is given next. This makes the theory of enrichment in an **fc**-multicategory run more smoothly (sometimes, at least) than that of enrichment in a bicategory, as we shall see towards the end of Section 3.

Definition 3 *Let C and D be $\mathcal{V}$ -enriched categories. A $\mathcal{V}$ -enriched functor $F : C \longrightarrow D$ consists of*

- *a function $F : C_0 \longrightarrow D_0$*

- *for each $a \in C_0$, a vertical 1-cell*

$$\begin{array}{c} C[a] \\ \downarrow F_a \\ D[F(a)] \end{array}$$

- *for each $a, b \in C_0$, a 2-cell*

$$\begin{array}{ccc} C[a] & \xrightarrow{C[a,b]} & C[b] \\ F_a \downarrow & \Downarrow F_{ab} & \downarrow F_b \\ D[F(a)] & \xrightarrow{D[F(a),F(b)]} & D[F(b)] \end{array}$$

such that the F_{ab} 's commute with the composition and identity 2-cells in C and D , in an evident sense.

With the obvious notion of composition of $\mathcal{V}$ -enriched functors, we obtain a category $\mathcal{V}\text{-}\mathbf{Cat}$ of $\mathcal{V}$ -enriched categories and functors.

Examples

1. Let $\mathcal{M}$ be a monoidal category and consider a category C enriched in the **fc**-multicategory $\Sigma\mathcal{M}$ (defined in Example 1(3)). There is only one possible choice for the $C[a]$'s, so the data for C consists of the set C_0 , the objects $C[a, b]$ of $\mathcal{M}$, and the maps

$$C[b, c] \otimes C[a, b] \longrightarrow C[a, c], \quad I \longrightarrow C[a, a].$$

Thus it turns out that a category enriched in $\Sigma\mathcal{M}$ is just a category enriched (in the usual sense) in $\mathcal{M}$. The same goes for enriched functors, so $(\Sigma\mathcal{M})\text{-}\mathbf{Cat}$ is isomorphic to the usual category of $\mathcal{M}$ -enriched categories and functors.

2. Let $\mathcal{M}$ be an (ordinary) multicategory. There is an obvious notion of category enriched in $\mathcal{M}$: that is, a set C_0 together with an object $C[a, b]$ of $\mathcal{M}$ for each $a, b \in C_0$ and arrows

$$C[a, b], C[b, c] \longrightarrow C[a, c], \quad \cdot \longrightarrow C[a, a]$$

(where $\cdot$ is the empty sequence), obeying suitable axioms. This is precisely the same thing as a category enriched in $\Sigma\mathcal{M}$.

3. If $\mathcal{B}$ is a bicategory then our $\mathcal{B}\text{-Cat}$ is isomorphic to the category of $\mathcal{B}$ -enriched categories defined in Walters [15].
4. Fix a topological space A . Then there is a bicategory $\Pi_2 A$, the *homotopy bicategory* of A , in which an object is a point of A , a 1-cell is a path in A , and a 2-cell is a homotopy class of path homotopies in A . For any 1-cell $\gamma : a \longrightarrow b$ there is an associated 1-cell $\gamma^* : b \longrightarrow a$ (that is, γ run backwards), and there are canonical 2-cells $1_b \longrightarrow \gamma \circ \gamma^*$ and $\gamma^* \circ \gamma \longrightarrow 1_a$.

Now suppose that A is nonempty and path-connected, and make a choice of a basepoint a_0 and for each $a \in A$ a path $\gamma_a : a_0 \longrightarrow a$. Then we obtain a category C enriched in $\Pi_2 A$, as follows:

- C_0 is the underlying set of A
- $C[a] = a$
- $C[a, b] = \gamma_b \circ \gamma_a^*$ (a path from a to b)
- composition $C[b, c] \circ C[a, b] \longrightarrow C[a, c]$ is the 2-cell

$$(\gamma_c \circ \gamma_b^*) \circ (\gamma_b \circ \gamma_a^*) \longrightarrow \gamma_c \circ \gamma_a^*$$

coming from the canonical 2-cell $\gamma_b^* \circ \gamma_b \longrightarrow 1_{a_0}$

- the identity 2-cell $1_a \longrightarrow C[a, a]$ is the canonical 2-cell $1_a \longrightarrow \gamma_a \circ \gamma_a^*$.

5. In the previous example, the bicategory $\Pi_2 A$ can be replaced by any bicategory $\mathcal{B}$ in which the underlying directed graph of objects and 1-cells is (nonempty and) connected, and every 1-cell has a left adjoint. (I thank the referee for alerting me to this.)
6. **Span-Cat** is equivalent to the comma category $(\text{ob} \downarrow \mathbf{Set})$, where $\text{ob} : \mathbf{Cat} \longrightarrow \mathbf{Set}$ is the objects functor. This means that a category enriched in **Span** consists of a category D , a set I , and a function $\text{ob}(D) \longrightarrow I$. To see why this is true, recall that a category C enriched in **Span** consists of
- a set C_0
 - for each $i \in C_0$, a set $C[i]$

- for each $i, j \in C_0$, a span

$$C[i] \xleftarrow{s_{ij}} C[i, j] \xrightarrow{t_{ij}} C[j]$$

- composition functions $C[j, k] \times_{C[j]} C[i, j] \longrightarrow C[i, k]$
- identity functions $C[i] \longrightarrow C[i, i]$,

all satisfying axioms. We can construct from C a category D with object-set $\coprod_{i \in C_0} C[i]$, arrow-set $\coprod_{i, j \in C_0} C[i, j]$, source and target maps given by the s_{ij} 's and t_{ij} 's, and composition and identity operations coming from those in C . By taking $I = C_0$ and the projection function $\text{ob}(D) \longrightarrow I$, we now have an object of $(\text{ob} \downarrow \mathbf{Set})$. A similar analysis of **Span**-enriched functors can be carried out, and we end up with a functor

$$\mathbf{Span-Cat} \longrightarrow (\text{ob} \downarrow \mathbf{Set}).$$

It is easy to see that this functor is an equivalence.

Let us briefly consider enriched categories with only one object. In the classical case of enrichment in a monoidal category $\mathcal{M}$, the category of one-object $\mathcal{M}$ -enriched categories is the category $\mathbf{Mon}(\mathcal{M})$ of monoids in $\mathcal{M}$. For an arbitrary **fc**-multicategory $\mathcal{V}$, we therefore define $\mathbf{Mon}(\mathcal{V})$ to be the full subcategory of $\mathcal{V}\text{-Cat}$ whose objects are $\mathcal{V}$ -enriched categories C with $|C_0| = 1$. Definitions 2 and 3 yield an explicit description of $\mathbf{Mon}(\mathcal{V})$.

Examples

1. If $\mathcal{M}$ is a monoidal category then $\mathbf{Mon}(\Sigma\mathcal{M})$ is the category of monoids in $\mathcal{M}$.
2. If $\mathcal{M}$ is a multicategory then an object of $\mathbf{Mon}(\Sigma\mathcal{M})$ consists of an object M of $\mathcal{M}$ together with maps

$$M, M \longrightarrow M, \quad \cdot \longrightarrow M$$

satisfying associativity and identity laws—in other words, a ‘monoid in $\mathcal{M}$ ’. A monoid in $\mathcal{M}$ is also the same thing as a multicategory map $\mathbf{1} \longrightarrow \mathcal{M}$, where $\mathbf{1}$ is the terminal multicategory.

3. If $\mathcal{B}$ is a bicategory then an object of $\mathbf{Mon}(\mathcal{B})$ is a monad in $\mathcal{B}$ in the sense of Street [14]: that is, it's an object X of $\mathcal{B}$ together with a 1-cell $t : X \longrightarrow X$ and 2-cells $\mu : t \circ t \longrightarrow t$, $\eta : \mathbf{1} \longrightarrow t$ satisfying the usual monad axioms. There are no maps $(X, t, \mu, \eta) \longrightarrow (X', t', \mu', \eta')$ in $\mathbf{Mon}(\mathcal{B})$ unless $X = X'$, and in this case such a map consists of a 2-cell $t \longrightarrow t'$ commuting with the μ 's and η 's. So $\mathbf{Mon}(\mathcal{B})$ is the category of monads and ‘strict monad maps’ in $\mathcal{B}$.

4. Let $\mathcal{B}$ be a 2-category. Associated to $\mathcal{B}$ is not only the **fc**-multicategory $\mathcal{V}$ of the previous example—which we now call $\mathcal{V}$ —but also two more **fc**-multicategories, $\mathcal{W}$ and $\mathcal{W}'$. Both $\mathcal{W}$ and $\mathcal{W}'$ are defined from double categories (see Example 1(1)), and in both cases an object is an object of $\mathcal{B}$, a vertical 1-cell is a 1-cell of $\mathcal{B}$, and a horizontal 1-cell is also a 1-cell of $\mathcal{B}$. In the case of $\mathcal{W}$, a 2-cell inside

$$\begin{array}{ccc} X & \xrightarrow{t} & Y \\ f \downarrow & & \downarrow g \\ X' & \xrightarrow{t'} & Y' \end{array}$$

is a 2-cell $t' \circ f \longrightarrow g \circ t$ in $\mathcal{B}$; in the case of $\mathcal{W}'$, it is a 2-cell $g \circ t \longrightarrow t' \circ f$ in $\mathcal{B}$. Composition and identities are defined in the obvious way.

Since $\mathcal{V}$, $\mathcal{W}$ and $\mathcal{W}'$ are identical when we ignore the vertical 1-cells, the objects of $\mathbf{Mon}(\mathcal{W})$ and $\mathbf{Mon}(\mathcal{W}')$ are the same as the objects of $\mathbf{Mon}(\mathcal{V})$; that is, they are monads in $\mathcal{B}$. But by using $\mathcal{W}$ or $\mathcal{W}'$ we obtain a more flexible notion of a ‘map of monads’ than we did in Example (3): a map in $\mathbf{Mon}(\mathcal{W})$ is what Street called a *monad functor* in [14], and a map in $\mathbf{Mon}(\mathcal{W}')$ is a *monad opfunctor*.

3 Bimodules

Bimodules have traditionally been discussed in the context of bicategories. Thus given a bicategory $\mathcal{B}$, one constructs a new bicategory $\mathbf{Bim}(\mathcal{B})$ whose 1-cells are bimodules in $\mathcal{B}$ (see e.g. [4]). The drawback is that this is only possible when $\mathcal{B}$ has certain properties concerning the existence and behaviour of local reflexive coequalizers.

Here we extend the **Bim** construction from bicategories to **fc**-multicategories, which allows us to drop the technical assumptions. In other words, we will construct an honest functor

$$\mathbf{Bim} : \mathbf{fc}\text{-}\mathbf{Multicat} \longrightarrow \mathbf{fc}\text{-}\mathbf{Multicat}.$$

(**fc**-**Multicat** is the category of (small) **fc**-multicategories, with maps defined in the obvious way.)

I would like to be able to, but at present cannot, place the **Bim** construction in a more abstract setting: as it stands it is somewhat *ad hoc*. In particular, the definition does not appear to generalize to T -multicategories for arbitrary T .

The theories of bimodules and enrichment interact in the following way: given an **fc**-multicategory $\mathcal{V}$, there is a canonically-defined functor

$$\mathcal{V}\text{-Cat} \longrightarrow \mathbf{Bim}(\mathcal{V})\text{-Cat}.$$

This is discussed at the end of the section, and provides lots of new examples of enriched categories.

We first have to define **Bim**. Let $\mathcal{V}$ be an **fc**-multicategory: then the **fc**-multicategory **Bim**($\mathcal{V}$) is defined as follows.

0-cells A 0-cell of **Bim**($\mathcal{V}$) is an **fc**-multicategory map $1 \longrightarrow \mathcal{V}$. That is, it is a 0-cell x of $\mathcal{V}$ together with a horizontal 1-cell $x \xrightarrow{t} x$ and 2-cells

$$\begin{array}{ccc} x & \xrightarrow{t} & x \xrightarrow{t} x \\ \downarrow 1 & \Downarrow \mu & \downarrow 1 \\ x & \xrightarrow{t} & x \end{array} \quad \begin{array}{ccc} x & \xRightarrow{\quad} & x \\ \downarrow 1 & \Downarrow \eta & \downarrow 1 \\ x & \xrightarrow{t} & x \end{array}$$

satisfying the usual axioms for a monad, $\mu \circ (\mu, 1_t) = \mu \circ (1_t, \mu)$ and $\mu \circ (\eta, 1_t) = 1_t = \mu \circ (1_t, \eta)$.

Horizontal 1-cells A horizontal 1-cell $(x, t, \eta, \mu) \longrightarrow (x', t', \eta', \mu')$ consists of a horizontal 1-cell $x \xrightarrow{m} x'$ in $\mathcal{V}$ together with 2-cells

$$\begin{array}{ccc} x & \xrightarrow{t} & x \xrightarrow{m} x' \\ \downarrow 1 & \Downarrow \theta & \downarrow 1 \\ x & \xrightarrow{m} & x' \end{array} \quad \begin{array}{ccc} x & \xrightarrow{m} & x' \xrightarrow{t'} x' \\ \downarrow 1 & \Downarrow \theta' & \downarrow 1 \\ x & \xrightarrow{m} & x' \end{array}$$

satisfying the usual module axioms $\theta \circ (\eta, 1_m) = 1_m$, $\theta \circ (\mu, 1_m) = \theta \circ (1_t, \theta)$, and dually for θ' , and the ‘commuting actions’ axiom $\theta' \circ (\theta, 1_{t'}) = \theta \circ (1_t, \theta')$.

Vertical 1-cells A vertical 1-cell $\begin{array}{c} (x, t, \eta, \mu) \\ \downarrow \\ (\hat{x}, \hat{t}, \hat{\eta}, \hat{\mu}) \end{array}$ in **Bim**($\mathcal{V}$) is a vertical 1-cell $\begin{array}{c} x \\ \downarrow f \\ \hat{x} \end{array}$ in $\mathcal{V}$ together with a 2-cell

$$\begin{array}{ccc} x & \xrightarrow{t} & x \\ f \downarrow & \Downarrow \omega & \downarrow f \\ \hat{x} & \xrightarrow{\hat{t}} & \hat{x} \end{array}$$

satisfying $\omega \circ \mu = \hat{\mu} \circ (\omega, \omega)$ and a similar equation for units.

2-cells A 2-cell

$$\begin{array}{ccccccc} t_0 & \xrightarrow{m_1} & t_1 & \xrightarrow{m_2} & \cdots & \xrightarrow{m_n} & t_n \\ f \downarrow & & & & \Downarrow & & \downarrow f' \\ t & \xrightarrow{\quad m \quad} & & & & & t' \end{array}$$

in $\mathbf{Bim}(\mathcal{V})$, where t stands for (x, t, η, μ) , m for (m, θ, θ') , f for (f, ω) , and so on, consists of a 2-cell

$$\begin{array}{ccccccc} x_0 & \xrightarrow{m_1} & x_1 & \xrightarrow{m_2} & \cdots & \xrightarrow{m_n} & x_n \\ f \downarrow & & & & \Downarrow \alpha & & \downarrow f' \\ x & \xrightarrow{\quad\quad\quad} & & & & & x' \\ & & & & & & m \end{array}$$

in $\mathcal{V}$, satisfying the ‘external equivariance’ axioms

$$\begin{aligned} \alpha \circ (\theta_1, 1_{m_2}, \dots, 1_{m_n}) &= \theta \circ (\omega, \alpha) \\ \alpha \circ (1_{m_1}, \dots, 1_{m_{n-1}}, \theta'_n) &= \theta' \circ (\alpha, \omega') \end{aligned}$$

and the ‘internal equivariance’ axioms

$$\begin{aligned} \alpha \circ (1_{m_1}, \dots, 1_{m_{i-2}}, \theta'_{i-1}, 1_{m_i}, 1_{m_{i+1}}, \dots, 1_{m_n}) &= \\ \alpha \circ (1_{m_1}, \dots, 1_{m_{i-2}}, 1_{m_{i-1}}, \theta_i, 1_{m_{i+1}}, \dots, 1_{m_n}) & \end{aligned}$$

for $2 \leq i \leq n$.

Composition and identities For both 2-cells and vertical 1-cells in $\mathbf{Bim}(\mathcal{V})$, composition is defined directly from the composition in $\mathcal{V}$, and similarly identities.

Incidentally, the category formed by the objects and vertical 1-cells of $\mathbf{Bim}(\mathcal{V})$ is $\mathbf{Mon}(\mathcal{V})$, the category of monads in $\mathcal{V}$ defined earlier.

We have now defined an **fc**-multicategory $\mathbf{Bim}(\mathcal{V})$ for each **fc**-multicategory $\mathcal{V}$, and it is clear how to do the same thing for maps of **fc**-multicategories, so that we have a functor

$$\mathbf{Bim} : \mathbf{fc}\text{-Multicat} \longrightarrow \mathbf{fc}\text{-Multicat}.$$

Again, we have been rather eccentric in naming the ‘bimodules construction’ after what it does to the horizontal 1-cells rather than the objects: perhaps we should call it the ‘monads construction’. We are, however, following the traditional terminology.

Examples

1. Let $\mathcal{B}$ be a bicategory satisfying the conditions on local reflexive coequalizers mentioned in the first paragraph of this section, so that it is possible to construct a bicategory $\mathbf{Bim}(\mathcal{B})$ in the traditional way. Let $\mathcal{V}$ be the **fc**-multicategory coming from $\mathcal{B}$. Then a 0-cell of $\mathbf{Bim}(\mathcal{V})$ is a monad in $\mathcal{B}$, a horizontal 1-cell $t \longrightarrow t'$ is a (t', t) -bimodule, and a 2-cell of the form

$$\begin{array}{ccccccc} t_0 & \xrightarrow{m_1} & t_1 & \xrightarrow{m_2} & \cdots & \xrightarrow{m_n} & t_n \\ 1 \downarrow & & & & \Downarrow & & \downarrow 1 \\ t_0 & \xrightarrow{\quad\quad\quad} & & & & & t_n \\ & & & & & & m \end{array}$$

is a map

$$m_n \otimes_{t_{n-1}} \cdots \otimes_{t_1} m_1 \longrightarrow m$$

of (t_n, t_0) -bimodules, i.e. a 2-cell in $\mathbf{Bim}(\mathcal{B})$. So if we discard the non-identity 1-cells of $\mathbf{Bim}(\mathcal{V})$, then the resulting **fc**-multicategory is precisely the **fc**-multicategory associated with the bicategory $\mathbf{Bim}(\mathcal{B})$.

2. $\mathbf{Bim}(\mathbf{Span}) = \mathbf{Prof}$, where $\mathbf{Span}$ is the **fc**-multicategory of sets, functions, spans, etc, and $\mathbf{Prof}$ is the **fc**-multicategory of categories, functors, profunctors, etc (Examples 1(5) and (6)).
3. $\mathbf{Bim}(\Sigma\mathbf{Ab}) = \mathbf{Bimod}$ (Example 1(7)). Here $\mathbf{Ab}$ is regarded as a monoidal category under the usual tensor and $\Sigma\mathbf{Ab}$ is as in Example 1(3); or equivalently $\mathbf{Ab}$ is regarded as a multicategory with the usual multilinear maps, and $\Sigma\mathbf{Ab}$ is as in Example 1(4).
4. Similarly, $\mathbf{Bim}(\Sigma\mathbf{Set}) = \mathbf{Action}$ (Example 1(8)), with cartesian product giving the monoidal category (or multicategory) structure on $\mathbf{Set}$.
5. It is possible to define an **fc**-multicategory $\mathbf{Span}(\mathcal{E}, T)$, for any appropriate monad T on a category $\mathcal{E}$, and then $\mathbf{Bim}(\mathbf{Span}(\mathcal{E}, T))$ is the **fc**-multicategory of T -multicategories and maps, profunctors, etc, between them. See [7] or [8] for details.

We now show how the bimodules construction produces new enriched categories from old.

Proposition 4 *For any **fc**-multicategory $\mathcal{V}$, there is a natural functor*

$$(\tilde{}) : \mathcal{V}\text{-}\mathbf{Cat} \longrightarrow \mathbf{Bim}(\mathcal{V})\text{-}\mathbf{Cat},$$

preserving object-sets.

Proof Take a $\mathcal{V}$ -enriched category C . We must define a $\mathbf{Bim}(\mathcal{V})$ -enriched category $\tilde{C}$ with object-set C_0 , and so, for instance, we must define an object $\tilde{C}[a]$ of $\mathbf{Bim}(\mathcal{V})$ for each $a \in C_0$. To do this we observe that $C[a]$ has a natural monad structure on it: that is, we put

$$\tilde{C}[a] = (C[a], C[a, a], \text{ids}_a, \text{comp}_{a, a, a}).$$

The rest of the construction is along similar lines; there is only one sensible way to proceed, and it is left to the reader. (An abstract account is in [8]). $\square$

Examples

1. Let C be a category enriched (in the usual sense) in the monoidal category $\mathbf{Ab}$ of abelian groups. Then the resulting $\mathbf{Bimod}$ -enriched category $\tilde{C}$ is as follows:
 - $\tilde{C}_0$ is the set of objects of C

- $\tilde{C}[a]$ is the ring $C[a, a]$, in which multiplication is given by composition in C
- $\tilde{C}[a, b]$ is the abelian group $C[a, b]$ acted on by $\tilde{C}[a] = C[a, a]$ on the right and by $\tilde{C}[b] = C[b, b]$ on the left, both actions being by composition in C
- composition and identities are as in C .

To illustrate the functoriality in the Proposition, take an **Ab**-enriched functor $F : C \longrightarrow D$. This induces a **Bimod**-enriched functor $\tilde{F} : \tilde{C} \longrightarrow \tilde{D}$ as follows:

- $\tilde{F} : C_0 \longrightarrow D_0$ is the object-function of F
- if $a \in C_0$ then $\tilde{F}_a$ is the ring homomorphism

$$\tilde{C}[a] = C[a, a] \longrightarrow D[F(a), F(a)] = \tilde{D}[\tilde{F}(a)]$$

induced by F

- if $a, b \in C_0$ then

$$\tilde{F}_{ab} : \tilde{C}[a, b] = C[a, b] \longrightarrow D[F(a), F(b)] = \tilde{D}[\tilde{F}(a), \tilde{F}(b)]$$

is defined by the action of F on morphisms $a \longrightarrow b$.

Note that in general, the ring homomorphism $\tilde{F}_a$ is not the identity; so the vertical 1-cells of **Bimod** get used in an essential way. This is the reason why the Proposition does not hold if we work throughout with bicategories rather than **fc**-multicategories: $(\tilde{})$ is defined on objects of $\mathcal{V}\text{-Cat}$, but cannot sensibly be defined on morphisms.

2. The non-additive version of (1) is that there is a canonical functor

$$\begin{array}{ccc} \mathbf{Cat} & \longrightarrow & \mathbf{Action-Cat} \\ C & \longmapsto & \tilde{C} \end{array}$$

which exists because, for instance, the set of endomorphisms on an object of a category is naturally a monoid.

3. In the previous example, part of the construction was to take $\tilde{C}[a]$ to be the monoid of all endomorphisms of a in C . However, we could just as well take only the automorphisms of a , and this would yield a different functor from **Cat** to **Action-Cat**.
4. Applying the Proposition to $\mathcal{V} = \mathbf{Span}$ and recalling Example 2(6), we obtain a functor

$$(\mathbf{ob} \downarrow \mathbf{Set}) \longrightarrow \mathbf{Prof-Cat}.$$

What this does on objects is as follows. Take a category D , a set I , and a function $\mathbf{ob}(D) \longrightarrow I$. Then in the resulting **Prof**-enriched category E

we have $E_0 = I$; $E[i]$ is the full subcategory of D whose objects are those lying over $i \in I$; and $E[i, j]$ is the profunctor

$$\begin{array}{ccc} E[i]^{\text{op}} \times E[j] & \longrightarrow & \mathbf{Set} \\ (d, d') & \longmapsto & D(d, d'). \end{array}$$

Composition and identities are defined as in D .

5. To get more examples of **Prof**-enriched categories we can modify the previous example, taking $E[i]$ to be *any* subcategory of D whose objects are all in the fibre over i . Here are two specific instances (each with a vague flavour of topological quantum field theory about them). In the first, E_0 is the set $\mathbb{N}$ of natural numbers, $E[n]$ is the category of n -dimensional Hilbert spaces (= complex inner product spaces) and isometries, and $E[m, n]$ sends (H, H') to the set of all linear maps $H \longrightarrow H'$. In the second, $E_0 = \mathbb{N}$ again, and we replace Hilbert spaces by differentiable manifolds, isometries by diffeomorphisms, and linear maps by differentiable maps.

Acknowledgements

I am grateful to Martin Hyland for his many words of wisdom, and for encouraging me to develop these ideas. I would also like to thank the organizers of CT99 at Coimbra for the opportunity to present this work and to discuss it with others. I thank the referee for a number of useful suggestions.

All but one of the diagrams in this document were generated using Paul Taylor's commutative diagrams package.

References

- [1] M. Batanin, Monoidal globular categories as a natural environment for the theory of weak n -categories, *Adv. Math.* 136 (1998), 39–103.
- [2] R. Borchers, Vertex algebras, e-print [q-alg/9706008](#), 1997.
- [3] A. Burroni, T -catégories, *Cahiers Top. Géom. Diff.* XII (1971), 215–321.
- [4] Aurelio Carboni, Stefano Kasangian, Robert Walters, An axiomatics for bicategories of modules, *J. Pure Appl. Algebra* 45 (1987), 127–141.
- [5] J. Lambek, Deductive systems and categories II, in: P. Hilton (ed.), *Category Theory, Homology Theory and their Applications I*, *Lecture Notes in Mathematics*, vol. 86, Springer, 1969, pp. 76–122.
- [6] Tom Leinster, General operads and multicategories, e-print [math.CT/9810053](#), 1997.
- [7] Tom Leinster, Generalized enrichment for categories and multicategories, e-print [math.CT/9901139](#), 1999.

- [8] Tom Leinster, Operads in higher-dimensional category theory, e-print [math.CT/0011106](#), Ph.D. thesis, University of Cambridge, 2000.
- [9] J. P. May, Definitions: operads, algebras and modules, in: Loday, Stasheff, Voronov (ed.), Operads: Proceedings of Renaissance Conferences, Contemporary Mathematics 202, AMS, 1997.
- [10] Craig T. Snyder, Equivalence of Borchers G -vertex algebras and axiomatic vertex algebras, e-print [math.QA/9904104](#), 1999.
- [11] Craig T. Snyder, Relaxed multi category structure of a global category of rings and modules, e-print [math.QA/9912075](#), 1999.
- [12] Y. Soibelman, Meromorphic tensor categories, e-print [q-alg/9709030](#), 1997.
- [13] Y. Soibelman, The meromorphic braided category arising in quantum affine algebras, Internat. Math. Res. Notices 19 (1999), 1067–1079.
- [14] Ross Street, The formal theory of monads, J. Pure Appl. Algebra 2 (1972), 149–168.
- [15] R. F. C. Walters, Sheaves and Cauchy-complete categories, Cahiers Top. Géom. Diff. XXII (1981), 283–286.